Index, Volume 51, 1958

Acalymma vittata, 81
Acarina, 259
Acaridiae, 767
Aceria
sheldoni, 657
tulipae, 68, 303
Acetals, as synergists in house fly sprays, 39
Acetates, myristyl and cetyl bromo- and chloro-, speed of action on resistant house flies, 880
Acethion and related compounds, toxicity, hydrolysis, and anticholinesterase activity of, 716
Achaea ablunaris, 637
Achaetoneura archippivora, 562
Achatina fulica, 390
Acheta assimilis, 866
Acids, bromo-, chloro-, and iodoacetic, speed of action on resistant house flies, 880
Acrospidia myrmicoides, 92
Acrostalagmus, probably *aphidium*, 184
Acrosternum
herbida, 636
vicina, 636
Acrylonitrile, dosages needed to kill rice weevil, 157
Adalia, 614
bipunctata, 81
Adelphocoris
lineolatus, 493, 729
rapidus, 493
Aedes
aegypti, 126, 187, 239, 345, 409, 499, 744, 780
atlanticus, 846
infirmatus, 846
sollicitans, 846
spp., 845
taeniorhynchus, 640, 846
triseriatus, 744
zoosophus, 744
Aedes, floodwater species of, 56
Aeolothrips duvali, 450
Aestivation of the alfalfa weevil, 506
Agathis stigmaterus, 164
Agonoderus comma, 81
Agriotes mancus, 690
Aircraft, jet, insect survival in, 239
Alabama argillacea, 237, 259, 633, 758, 783
Aldrin
as a fumigant against house flies, 638
as a seed treatment against wireworms, 294
dips to reduce weevil damage to pine seedlings, 916
distribution of in soils, 381
Aldrin, conversion to dieldrin,
in soil, 1
on plants, 8
Aldrin, for control of
alfalfa snout beetle, 683
carrot rust fly, 166
clover leaf weevil, 195
European chafer, 766
Nysius raphanus, 251
onion maggot, 884
paddy borer, 571
plum curculio, 131
serpentine leaf miner, 358
sugarcane beetle, 631
weevils on crimson clover, 459
western subterranean termite, 209
wireworms, 464, 691
Aldrin
granules, performance of broadcaster with, 280
oral chemotherapeutic activity of in mice, 127
percentage recovered from treated soil, 201
persistence and transformation of in soils, 214
residues of in carrots, 337
residues on alfalfa, 222
Aldrin, toxicity of to
army cutworm, 726
brown-banded cockroach, 619
Ceratitis capitata, 235
pink bollworm adults, 578
young cabbage and muskmelon plants, 534
Alfalfa
insect-free, seed production and growth characteristics of, 729
reaction of to spotted alfalfa aphid, 601
resistant to meadow spittlebug, 219
Alkoxybenzyl esters of chrysanthemumic acid, toxicity of in house fly sprays, 810
Allethrin
cis isomers of, mixed with piperonyl butoxide, 675
four *cis* isomers of, 181
toxicity of in house fly sprays, 810
with acetal synergists in house fly sprays, 39
Allethrin
cis and *trans* fractions of, intensity of synergism of their mixtures with piperonyl butoxide, 76
relative toxicity to house flies, 76
Allotropa citri, 90
Alphitobius diaperinus, 112
Altica ambiens, 296
Amblyomma
americanum, 425
maculatum, 499
Amblyseius sp., 366
Am. Cyanamid 4124 (see also dicapthon)
for control of bed bug, 241
toxicity of to honey bees, 103
Am. Cyanamid 12008, for control of
Aceria tulipae, 70
greenbug, 723
green peach aphid, 139
spotted alfalfa aphid, 122
thrips on cotton, 160
two-spotted spider mite, 736
Am. Cyanamid 12008
metabolism of by some soil organisms, 59
toxicity of to honey bees, 103
translocation and persistence of in cacao foliage, 773
Am. Cyanamid 12009, for control of
Aceria tulipae, 70
thrips on cotton, 160
Am. Cyanamid 12009
toxicity of to honey bees, 103
translocation and persistence of in cacao foliage, 773
Am. Cyanamid 12880,
as a chemotherapeutic agent, 126
bioassay of blood from cattle treated with, 861
Am. Cyanamid 12880, for control of
balsam gall midge, 400
cattle grubs, 445, 583, 793, 876
cattle lice, 876
lone star tick, 425
screw-worm, 425
stable fly, 425
Am. Cyanamid 12880, radioactive, in mosquitoes, 640
Am. Cyanamid 18706, for control of two-spotted spider mite, 736
Ammonium carbonate to attract walnut husk fly, 687
Amorbia essigana, 296
Amphimallon majalis, 486, 765, 869, 912
Anagyrus
antoninae, 363
diversicornis, 365
graminicolens, 365
kivuensis, 92
pseudococci, 90
Analyses
microbial, from insecticide-treated soils, 200
physical, chemical, and microbial of Chehalis and Olympic silt loam soils, 204
probit, on a digital computer, 738
Anaphothrips zeae, 450
Anarhopus sydneyensis, 92
Anasa tristis, 169
Anastatus blattidarum, 608
Androlaelaps sp., 366
Anoetus sp., 461
Anopheles
barberi, 744
quadrimaculatus, 240, 345, 640
Anthonomus
grandis, 41, 93, 100, 114, 236, 288, 502, 508, 669, 719, 745, 783, 872
sp., 636
Anthrenus flavipes, 248
Anticarsia gemmatalis, 636
Anticholinesterase agents, recovery of from cacao beans treated with,
BHC, 778
Chipman R-6199 (free base), 777
Chipman R-6200, 778
demeton, 777
dimefox, 778
Isolan, 777
Thimet (Oxygen analog sulfone), 777
Antonina
graminis, 363
indica, 363
phragmitis, 365
purpurea, 365
Anuraphis roseus, 354
Anystis sp., 366
Aonidiella aurantii, 193, 232, 254, 474, 764 911
Apanteles
angaleti, 164
chelonis, 390
Aphelinus
nigritus, 553
semiflavus, 374
Aphicides, selective, favoring natural enemies of the spotted alfalfa aphid, 374
Aphidius testaceipes, 553
Aphids

control of on cotton with systemics, 699
effect of low temperatures on survival of, 144
populations on field beans in Washington, 178
provide substitute sugar source for honey bees, 287
Aphis
abbreviata, 144, 414
crataequifoliae, 179
fabae, 735
gossypii, 41, 45, 236, 636, 872
helichrysi, 226
knowltoni, 179
phaceliae, 179
pomi, 357, 422
spiraecola, 179
Aphodius distinctus, 81
Aphytis
chrysomphali, 476
lingnanensis, 476
Apis mellifera, 46, 64, 103, 146, 465, 494, 523, 784
Aramite, for control of
chicken body louse, 230
citrus bud mite, 659
cyclamen mite, 432
mites attacking eye gnat cultures, 462
two-spotted spider mite, 353, 356, 711, 714
Aramite, relative toxicity of to five species of tetranychid mites, 521
Aramite, resistance to by
Tetranychus atlanticus, 286
Tetranychus pacificus, 286
Archips
argyrospila, 598
rosana, 89
Archytas marmoratus, 562
Argyrotaenia
citrana, 89, 596
velutinana, 81, 357, 378, 454
Arhyssus parviconis, 636
Aroclor, for control of house flies, 20
Arthrochlamps plicata, 81
Aspidiotiphagus citrinus, 115
Artemia salina, use of for bioassay, 780
Asca
bicornis, 770
sphidioides, 770
Aster-yellows virus infection and transmission, 512
Astymachus sp., 366
AT-4, for control of horn flies and stable flies, 73
Athaumastus haematicus, 636
Attaclay, as a carrier for insecticides against wireworms, 463
Attagenus piceus, 248, 249, 887
Attractant, insect, effect of *cis-trans* isomerism on the potency of, 921
Autoserica castanea, 547
Attractants
for walnut husk fly, 689
glycine-sodium hydroxide solution for walnut husk fly, 532

Bacillus thuringiensis, 390
for control of codling moth, 422
susceptibility of some insects to, 296
Baits
attractants for walnut husk fly, 689
against house flies, 19
Dipterex against house flies, 19, 798
DDVP against house flies, 19
DDT against confused flour beetle, 301
for control of *Fannia canicularis*, 611
for control of *Leptocorixa acuta*, 405
lures for Mediterranean fruit fly in Florida, 679
malathion against house flies, 19
Pival against confused flour beetle, 301
rodenticidal, resistance of to *Tribolium confusum*, 299
synergized pyrethrins and allethrin to control house flies, 798
Balaustium sp., 366
Barichneumon, 652
Bathyplectes curculionis, 298
Bayer 13,242, systemic activity against the bed bug, 499
Bayer 17,364, systemic activity against the bed bug, 499
Bayer 18,613, systemic activity against the bed bug, 499
Bayer 19,596, systemic activity against the bed bug, 499
Bayer 19,639
for control of boll weevil, 100
citrus whitefly, 552
cotton aphid, 100
iris borer, 550
onion maggot, 885
spotted alfalfa aphid, 122, 457
thrips on cotton, 160
two-spotted spider mite, 736
Bayer 19,639
effect of seed treatments on cotton, 495, 697
resistance of citrus red mite to, 233
tests with, as cottonseed treatments, 872
translocation and persistence of in cacao foliage, 773
translocation of in pineapple plants, 905
Bayer 19,994, systemic activity against the bed bug, 499
Bayer 21/199, fate of in the white rat, 204
Bayer 21/199, for control of
cattle grubs, 359, 545, 583, 793, 876
cattle lice, 876
horn flies, 559, 746, 795
lone star tick, 425
onion maggot, 886
screw-worm, 408, 425
stable fly, 425
Bayer 21/199
radioactive, in mosquitoes, 640
resistance of house fly to, 497
Bayer 23,129, for control of
lone star tick, 425
screw-worm, 425
spotted alfalfa aphid, 123, 457
stable fly, 425
two-spotted spider mite, 736
Bayer 23,129, translocation and persistence of in cacao foliage, 773
Bayer 23,453, for control of
lone star tick, 425
screw-worm, 425
stable fly, 425
Bayer 25,141, for control of boll weevil, 785
Beauveria bassiana, 391
Behavior, wireworm, in response to chemical seed treatments, 291
Bergoldia clistorhabdion, 454
BHC dips to reduce weevil damage to pine seedlings, 916
BHC, effect of on
parasites of oriental fruit moth, 85
soft scales on walnut, 429
BHC, for control of
alfalfa snout beetle, 683
apple grain aphid, 355
boll weevil, 784
cotton leafworm, 784
Epilachna vigintioctopunctata, 749
European chafer, 766
greenbug, 723
peach tree borer, 558
root collar borer in yellow-poplar, 252
rosy apple aphid, 355
serpentine leaf miner, 358
walnut scale, 528
BHC, polyfluoro analogs of, 413
BHC, residues in treated soil, 187
BHC, toxicity of
in milk to house flies, 560
to pink bollworm adults, 578
to *Therioaphis maculata* and *Hippodamia*, 377
Bioassay
for testing effectiveness of malathion and lindane against wheat insects, 843
of carrots for aldrin residue, 337
to determine concentration of Bayer 19,639 in pineapple plants, 905
to determine insecticide residues in soils, 186
to determine persistence of aldrin and heptachlor residues on alfalfa, 222
to determine residue of dieldrin on alfalfa, 685
to study decline of soil insecticides, 200
use of brine shrimp as a test organism, 780
with stable fly, of blood from cattle treated with Am. Cyanamid 12880, 861
Biological Control
agents introduced into Indonesia, 390
characteristics of Ichneumonidae in relation to, 650
Rhodes-grass scale in Texas, 363
Biological effectiveness of
insect screens, 864
Biology of
Aphis helichrysi, 226
brine shrimp, 781
brown-banded cockroach, 608
carrot weevil in Illinois, 37
cherry fruit flies on Vancouver Island, 592
citrus cutworm, 653
Euzophora ostricolorella, 251
Fiorinia externa, 405
McDaniel mite in Utah, 588
purple-backed cabbageworm, 281
seed-corn maggot, 704
vetch bruchid, 401
Bio 1240, resistance of citrus red mite to, 232
Black Leaf 40, for control of chicken body louse, 230
Blatta orientalis, 866
Blattella germanica, 211, 239, 573, 608, 866, 867
Blattisocius keegani, 366
Blepharigena spinosula, 562
Blepyrus saccharicola, 92
Blueberries, low-bush, pollination, of 544
Boopedon sp., 465
Bothriocraera bicolor, 92
Boucekiella antoninae, 365
Brachymeria ovata abiesae, 764
Brachyrhinus ligustici, 682
Bracon
brevicornis, 164
xanthonotus, 764
Brevicoryne brassicae, 553, 626
Brevipalpus phoenicis, 636
Bruchidae, 636
Bruchophagus

gibbus, 409
kolobovae, 409
Bruchus brachialis, 401
Brumus suturalis, 92
Bryobia praetiosa, 767
Buccalatrix thurberiella, 296
Bulan, for control of serpentine leaf miner, 358
Butoxy polypropylene glycol, for control of
horn flies, 73, 149
stable flies, 73, 149
Butoxy polypropylene glycol in treatments against biting flies on cattle, 620

Calacoris lineolatus, 636
Calcium arsenate, for control of
bollworm, 42, 646
boll weevil, 42
serpentine leaf miner, 358
Callisto geminatella, 81
Callitroga
hominivorax, 407, 425
macellaria, 407
Capitophorus
braggi, 179
cardiunus, 179
hippophaes, 179
shepheridiae, 179
Captan, in spray combinations against apple insects, 378
Carabid beetles damage Douglas-fir seed, 542
Carbamoyl alkyl phosphorodithioates as chemotherapeutic agents, 445
Carbon disulfide, dosages needed to kill rice weevil, 157
Carpocapsa pomonella, 81, 354, 378, 422, 542, 547, 672, 693
Casinaria sp., 390
Cat-facing in peaches, 27
Cattle, beef
effect of biting fly control on weight gain in, 275
weight responses of in relation to control of flies, 72
Cattle, control of biting flies on, 149
Cattle, dairy, relationship of stable flies to milk production, 269
Cephus cinctus, 804
Cerapterocerus mirabilis, 365
Ceratitus capitata, 234, 679, 921
Ceratocapsus nigellus, injury to peaches, 28
Ceratocystis ulmi, 256
Cerotoma trifurcata, 81
Certacopsis sp., 636
Chaetogaedia sp., near *analis*, 562
Chaetopsis fulvifrons, 81
Chalcids, seed, reared from birdsfoot trefoil, 409
Chalcodermus sp., 636
Cheiloneurus
elegans, 365
sp., 366
Chelonus
heliopae, 164
munakatae, 390
n. sp., 164
Chelycoris sp., 636
Chelymorpha sp., 636
Chermes piceae, 540
Chemotherapeutic agents
phosphorodithioates, carbamoylalkyl, 126
Cheyletia wellsi, 366
Cheyletidae, 366
Chilocorus
angolensis, 92
bivulnerus, 81
spp., 479
Chilo suppressalis, 391, 571
Chipman R-6199, for control of
balsam gall midge, 400
boll weevil, 785
codling moth, 379
desert spider mite, 646
European red mite, 714
red-banded leaf roller, 379
Scolytus multistriatus, 789
two-spotted spider mite, 714
Chipman R-6199
resistance of citrus red mite to, 233
translocation and persistence of in cacao foliage, 773
Chipman R-6200
for control of citrus mealybug, 779
translocation and persistence of in cacao foliage, 775
Chirothrips
falsus, 450
mexicanus, 450
Chlorbenside, for control of European red mite, 713
Chlorbenside, toxicity of to honey bees, 103
Chlordane, for control of
Aceria tulipae, 70
alfalfa snout beetle, 683
bed bug, 242
black carpet beetle, 249
carrot rust fly, 166
cherry fruit flies, 594
chicken body louse, 230
European chafer, 766
horn fly, 795
onion maggot, 884
plum curculio, 132
serpentine leaf miner, 358
sugarcane beetle, 632
tropical rat mite, 413
wireworms, 691
Chlordane
oral chemotherapeutic activity of in mice, 127
resistance of house fly to, 497
Chlordane, toxicity of to
army cutworm, 726
brown-banded cockroach, 610
Ceratitis capitata, 235
melon fly, 236
young cabbage and muskmelon plants, 534
Chlordane-treated soil effective against western subterranean termite, 207
Chlorella pyrenoidosa, 59
Chlorobenzilate, for control of
Aceria tulipae, 70
chicken mite, 911
citrus bud mite, 659
European red mite, 714
mites attacking eye gnat cultures, 462
two-spotted spider mite, 714
Chlorogenus callistephi vulgaris, 512
Chloropicrin, dosages needed to kill rice weevil, 157
Chlorthion, for control of
Aceria tulipae, 70
bed bug, 241
cabbage leaf-feeding lepidoptera, 443
clover leaf weevil, 195
corn earworm, 823
greenbug, 722
green peach aphid, 139
holly bud moth, 88
Mexican bean beetle, 554
Nysius raphanus, 251
onion maggot, 885
serpentine leaf miner, 358
sugarcane beetle, 633
thrips on cabbage, 395
two-spotted spider mite, 736
western tussock moth, 764
Chlorthion
hydrolysis of the alkyl-phosphate bond by rats, cockroaches, and alkali, 800
oral chemotherapeutic activity of in mice, 127
Cholinesterase
activity of bovine red-blood-cell, measurement of, 326
brain, inhibition and recovery of activity in house flies, 750
studies, ET-57 depressed enzyme activity, 121
Choristoneura fumiferana, 264
Chorizagrotis auxiliaris, 726
Chromaphis juglandicola, 427, 527
Chrysodina sp., 636
Chrysomya
megacephala, 147
sp., 147
Chrysopa, 651
Chrysopa
plorabunda group, 366, 613
spp., 90, 713, 757
Chrysoplatycerus splendens, 92
Chrysops univittata, 845
Chymomyza amoena, 734
Cimex
hemipterus, 241
lectularius, 241, 499
Circulifer tenellus, curly top-infective, effect of on watermelon plants, 434
Citrus
mealybugs attacking in California, 90
transpiration of, affected by spray oil, 317
Clepsis peritana, 596
Cnephia pecuarum, 845
Coccinella
californica, 90
novemnotata, 614
transversoguttata, 614
Coccidoxenus sp., 365
Coccophagus
gurneyi, 92
scutellaris, 247
Coccus hesperidum, 480
Coccygomimus sanguinipes, 764
Coeloides dendroctoni, 604
Coelomera sp., 636
Colaspis sp., 636
Coleomegilla
fuscilabris, 81
maculata, 759
octodecimpustuluta, 637
Colias philodice eurytheme, 562
Colladonus clitellarius, 81
Collembola, 252, 767
Collops bipunctata, 613
Compound K-8025, for control of garden tortrix, 597
Compounds
Ammonia-producing, to attract walnut husk fly, 688
biologically active, effect of on house fly ovaries, 384
Compounds
neuropharmacological, ionization and toxicity to insects, 169
neurotoxic toxicity of to the mouse, house fly and cockroach, 171
Compounds, synthetic,
screening tests of 111 for toxicity to house flies, 791

esters of chrysanthemumic acid toxic to house flies, 791
Compounds used in house fly studies,
carbamate, 750
organophosphate, 750
sulfamate, 750
Conocephalus
saltator, 147
strictus, 345
Conotelus mexicanus, 147
Conotrachelus nenuphar, 81, 131, 330, 357
Copper oxychloride sulfate, toxicity of to honey bees, 103
Corn, damage to by southern green stink bug, 471
CP 7769, for control of cotton leafworm, 259
Cotton, Upland, genotypic reaction to boll weevil attack in, 508
Cotton
resistance to boll weevil in mixed populations of resistant and susceptible plants, 502
seedling, Collembola damage to, 252
Crambus calliginosellus damaging carrots, 402
Cremastus chinensis, 390
Crimson clover seed, influence of honey bees on production of, 523
Crioceris duodecimpunctata, 81
Cryolite, for control of
citrus cutworm, 656
garden tortrix, 597
western tussock moth, 764
Cryptolaemus montrouzieri, 90
Culex
gelidus, 109
pipiens, 345
quinquefasciatus, 386, 744
tarsalis, resistance to DDT, 56
tritaeniorhynchus, 109
Culicoides spp., 845
Cultures, bacterial, for control of tobacco insects, 560
Cunaxa sp., 366
Cunaxidae, 771
Cyclocephala borealis, 547
Cyclohexanes, fluorinated, toxicities of to house flies, 413
Cycloneda
munda, 81
sanguinea, 637

Dacus
cucurbitae, 140, 147, 234
dorsalis, 147, 234
Danaus plexippus, attacking sour cherries, 298
DDD, (TDE) for control of codling moth, 542
DDT,
compounds related to evaluated as synergists of, 647
distribution of in soils, 381
DDT, effect of on
California red scale, 483
insect predators on alfalfa, 615
parasites of oriental fruit moth, 86
reproduction of house flies, 579
DDT, for control of
alfalfa pests, 730
black carpet beetle, 249
boll weevil, 42, 237, 784
bollworm, 42, 237, 646
cabbage leaf-feeding lepidoptera, 443
cabbage looper, 113, 237
carrot rust fly, 166
chicken body louse, 229
citrus cutworm, 656
codling moth, 357, 378, 422, 542
Colorado potato beetle, 829
corn earworm, 25, 823
Epilachna vigintioctopunctata, 749
European corn borer, 133
Fiorinia externa, 406
garden tortrix, 597
greenbug, 722
holly bud moth, 88
house flies, 20, 809
imported cabbageworm, 254
Japanese beetle, 247
lesser mealworm, 112
Lithocolletis sp., 423
meadow spittlebug, 493
onion maggot, 887
peach tree borer, 558
pink bollworm, 673
purple-backed cabbageworm, 284
Pyroderces rileyi, 24
red-banded leaf roller, 378
rice weevil, 25
root collar borer in yellow-poplar, 252
Scolytus multistriatus, 789
seed-corn maggot, 919
serpentine leaf miner, 358
six-spotted leafhopper, 903
smaller European elm bark beetle, 470
thrips on cabbage, 395
western subterranean termite, 207
western tussock moth, 764
DDT for testing insecticide deposits
with newly hatched codling moth larvae, 547
in bait against confused flour beetle, 301
in oil and in acetone, toxicity to resistant house flies, 914
oral chemotherapeutic activity of in mice, 127
radioactive, in mosquitoes, 640
rates of application to avoid loss of birds and fish, 741
residues of in treated soil, 187
residues of on vegetables, 818
resistance, an attempt to reduce in body lice, 665
susceptibility of to brine shrimp, 780
tests against house fly, 452
use of, causes increase of soft scales on walnut, 427
on vegetables, 307
DDT, resistance of
codling moth to, 693
Colorado potato beetle to, 828
house fly to, 497
potato flea beetle to, 823
stable flies to in Norway, 599
DDT, toxicity of compared with diphenylmethane against the confused flour beetle, 388
DDT, toxicity of to
brown-banded cockroach, 610
buckthorn aphid, 414
foxglove aphid, 414
goldfish, 540
green peach aphid, 414
honey bees, 103
pink bollworm adults, 568
potato aphid, 414
Scolytus multistriatus, 254
DDVP, as a fumigant against house flies, 638
DDVP, for control of
Fannia canicularis, 611
house flies, 20
Mediterranean fruit fly, 679
two-spotted spider mite, 736
DDVP, toxicity of to honey bees, 103
Dehydrogenase systems from fat body of *Blattella germanica*, 867
Demeton, for control of
Aceria tulipae, 70
balsam gall midge, 400
cabbage leaf-feeding lepidoptera, 443
citrus whitefly, 552
corn earworm, 259
European red mite, 255
greenbug, 723
green peach aphid, 139
leaf curl plum aphid, 228
Mexican bean beetle, 259
Nysius raphanus, 251
potato leaf hopper, 259, 336
red-headed pine sawfly, 400
resistant spotted alfalfa aphids, 313
Scolytus multistriatus, 789
serpentine leaf miner, 358
six-spotted leafhopper, 903
soft scales on walnut, 429
thrips, 160, 395
two-spotted spider mite, 711, 714, 736
walnut scale, 528
Demeton
oral chemotherapeutic activity of in mice, 127
residues of on cabbage and celery, 175
translocation and persistence of in cacao foliage, 773
Demeton, resistance to by
citrus red mite, 232
Tetranychus atlanticus, 285
Tetranychus pacificus, 285
Demeton, toxicity of to internal parasites of spotted alfalfa aphid, 375
Dendroctonus
brevicomis, 264
engelmanni, effects of woodpeckers on, 603
Deraecoris
historio, 81
quercicola, injury to peaches, 28
Dermacentor variabilis, 344
Dermanyssus gallinae, 158, 911
Dermatobia hominis, 724
Derris, for control of cherry fruit flies, 594
Desmometopa sp., 147
Diabrotica
speciosa, 636
undecimpunctata howardi, 81
Diachlorus ferrugatus, 845
Dialeurodes citri, 552
Diatraea saccharalis, 163, 562
Diazinon, effect of on
cocooning codling moth larvae, 672
reproduction of house flies, 579
Diazinon, for control of
Aceria tulipae, 70
bagworm, 368
bed bug, 241
black carpet beetle, 249
cherry fruit flies, 595
chicken mite, 911
clover leaf weevil, 196
codling moth, 422
corn earworm, 259
desert spider mite, 646
garden tortrix, 597
Fannia canicularis, 611
holly bud moth, 88
horn flies, 559
house flies, 20, 908
leaf curl plum aphid, 228
Mexican bean beetle, 259, 554
Nysius raphanus, 251
onion maggot, 885
potato leafhopper, 259
serpentine leaf miner, 358

western subterranean termite, 209
western tussock moth, 764
Diazinon
hydrolysis of the alkyl-phosphate bond by rats, cockroaches, and alkali, 800
oral chemotherapeutic activity of in mice, 127
Diazinon, resistance to of
citrus red mite, 233
house fly to, 497, 546, 616
Dibrachys cavus, 764
Dibrom for control of
iris borer, 550
two-spotted spider mite, 736
Dicapthon (see also Am. Cyanamid 4124) for control of
chicken mite, 911
citrus cutworm, 656
western tussock moth, 764
Dicapthon
hydrolysis of the alkyl-phosphate bond by rats, cockroaches, and alkali, 800
resistance of house fly to, 497
Dichelops melacanthus, 636
Dieldrin
as a seed treatment against wireworms, 294
as a permanent mothproofing agent, 887
conversion of aldrin to in soil, 1
detection of by potato flea beetle, 825
dips to reduce weevil damage to pine seedlings, 916
effect on insect predators on alfalfa, 615
from converted aldrin in carrots, 340
susceptibility of to brine shrimp, 780
Dieldrin, for control of
Aceria tulipae, 70
alfalfa snout beetle, 682
bagworm, 368
bed bug, 241
black carpet beetle, 249
boll weevil, 237, 490
bollworm, 238, 490
cabbage leaf-feeding lepidoptera, 443
carrot rust fly, 166
clover leaf weevil, 196
Colorado potato beetle, 829
cotton aphids, 238
cotton fleahopper, 490
European chafer, 766
Fuller rose beetle, 284
meadow spittlebug, 493
northern fowl mite, 188
Nysius raphanus, 251
onion maggot, 884
pink bollworm, 238
plum curculio, 131
rice paddy borer, 571
seed-corn maggot, 919
serpentine leaf miner, 358
sheep ked in Oregon, 585
smaller European elm bark beetle, 470
sugarcane beetle, 631
thrips, 395, 489
two-spotted mite, 238
weevils on crimson clover, 459
wireworms, 691
Dieldrin
oral chemotherapeutic activity of in mice, 127
percentage recovered from treated soils, 201
persistence and transformation of in soils, 214
residues in treated soil, 187
residues of on alfalfa, 392
resistance of house fly to, 497
Dieldrin, toxicity of to
army cutworm, 726
boll weevil, 670
Ceratitis capitata, 235
pink bollworm adults, 568
Scolytus multistriatus, 257
young cabbage and muskmelon plants, 534
Dieldrin-treated soil, effect of on
adults of European chafer, 486
western subterranean termite, 207
Diethyltoluamide
in treatments against biting flies on cattle, 620
ultraviolet determination of, 694
Diets
media for pink bollworm, 309
larval, for rearing the boll weevil, 288
Digamasellus angulosus, 770
Dilan, for control of
alfalfa snout beetle, 683
citrus cutworm, 656
garden tortrix, 597
serpentine leaf miner, 358
western tussock moth, 764
Dilan, toxicity of in soil to *Ceratitis capitata*, 235
Dimefox biological and chemical oxidation of, 49
Dimefox for control of
balsam hall midge, 400
citrus mealybug, 779
Scolytus multistriatus, 789
Dimefox, not oxidized in *Chlorella* cultures, 59
Dimite, for control of mites attacking eye gnat cultures, 462
Dimethoate (see Am. Cyanamid 12880)
Dinitrobutylphenol, for control of leaf curl plum aphid, 228
Dinitrocresol, for control of European red mite, 713
Dinitro phenol, for control of leaf curl plum aphid, 228
Dinitro 289, for control of soft scales on walnut, 430
Diphacinone, ineffective against confused flour beetle, 302
Diphenylmethane, derivatives, relation of structure to toxicity to confused flour beetle, 385
Dips, preplanting, reduce weevil damage to pine seedlings, 915
Dipterex, for control of
bed bug, 241
boll weevil, 42, 784
bollworm, 42
cabbage leaf-feeding lepidoptera, 443
codling moth, 422
house flies, 20, 453, 908
Mexican bean beetle, 554
onion maggot, 885
serpentine leaf miner, 358
thrips on cabbage, 395
two-spotted spider mite, 736
western tussock moth, 764
Dipterex
oral chemotherapeutic activity of in mice, 127
resistance of house fly to, 497, 616
toxicity of to honey bees, 103
Disonycha pennsylvanica, 81
Dissosteira carolina, 345
DNBP, for control of
apple grain aphid, 355
rosy apple aphid, 355
DN-40, for control of
apple grain aphid, 355
rosy apple aphid, 355
Douglas-fir seed, damage to by carabid beetles, 542
Dowco 109, for control of
cattle grubs, 583, 876
cattle lice, 876
Dow ET-15, for control of
boll weevil, 42
bollworm, 42
house flies, 20
Dow ET-15, toxicity of to honey bees, 102
Dow ET-57 as a fumigant against
American cockroach, 409
confused flour beetle, 409
house fly, 409, 638
yellow-fever mosquito, 409
Dow ET-57, for control of
bed bug, 241
cattle grubs, 119, 545, 583, 783, 876
cattle lice, 867
chicken body louse, 189, 230
chicken shaft louse, 189, 559
fluff louse, 189
house fly, 361, 908
horn flies, 559
lone star tick, 425
northern fowl mite, 189
onion maggot, 884
screw-worm, 425
serpentine leaf miner, 358
stable fly, 425
sticktight flea, 189
wing louse, 189
Dow ET-57
effect of treatment time on control of cattle grubs, 808
fumigation properties of, 408
hydrolysis of the alkylphosphate bond by rats, cockroaches, and alkali, 800
not oxidized in *Chlorella* cultures, 59
oral chemotherapeutic activity of in mice, 127
resistance of house fly to, 497
toxicity of to honey bees, 103
Draeculacephala mollipes, 81
Drepanaphis granovskyi, 179
Drosophila
affinis, 733
busckii, 733
duncani, 734
funebris, 733
immigrans, 733
melanogaster, 12, 81, 179, 222, 337, 685, 733
putrida, 733
quinaria, 733
repleta, 734
robusta, 733
tripunctata, 733
transversa, 734
Dysdercus pallidus, 633
Dysinicoccus obesus, 555

Echidnophaga gallinacea, 188
Ectoparasites on Alabama mammals, 260
Edessa
meditabunda, 637
sp., 636
Empoasca fabae, 220, 258, 336, 729
Empusa
aphidis, 185
no sp., 184
sphaerosperma, 185
Endrin, as a seed treatment against wireworms, 294
Endrin, effect of
climatic factors on toxicity of to boll weevil, 670

on cocooning colding moth larvae, 672
Endrin, for control of
Aceria tulipae, 70
alfalfa snout beetle, 683
boll weevil, 42, 95, 784
bollworm, 42, 237, 646
cabbage leaf-feeding lepidoptera, 443
cabbage looper, 237
clover leaf weevil, 195
corn earworm, 25
cotton leafworm, 237, 259, 784
cyclamen mite, 431
Epilachna vigintioctopunctata, 749
European chafer, 766
meadow spittlebug, 493
Nysius raphanus, 251
onion maggot, 884
paddy borer, 411, 571
pink bollworm, 673
Pyroderces rileyi, 25
rice weevil, 25
seed-corn maggot, 919
serpentine leaf miner, 358
salt-marsh caterpillar, 237
sugarcane borer, 562
thrips on cabbage, 395
Endrin, residues of on vegetables, 818
Endrin, toxicity of to
army cutworm, 726
Ceratitis capitata, 235
melon fly, 236
pink bollworm adults, 578
young cabbage and muskmelon plants, 534
Enoplopactus heterothorax, 636
Entomobrya unostrigata, 252
Entomologist, or forester, 264
Entomophagus species, role of in natural control of insect populations, 474
Entomophthora
exitialis, 341
oronata, 341
virulenta, 341
ENT 21478, as a lure for Mediterranean fruit fly, 679
Eomenacanthus stramineus, 188
Ephestia
elutella, 417
figulilella, 596
Epilachna
varivestis, 258, 345, 553, 652, 735
vigintioctopunctata, 749
Epitrix cucumeris, feeding behavior, 823
EPN, effect of on parasites of oriental fruit moth, 83
EPN, for control of
boll weevil, 42, 784
bollworm, 42
corn earworm, 823
cyclamen mite, 432
house flies, 20, 908
plum curculio, 132
serpentine leaf miner, 358
wireworms, 691
EPN, resistance of citrus red mite to, 233
Equipment
automation system for insectaries, 109
black-light traps used to attract hornworm moths, 79
broadcaster for applying granulated insecticides, 278
device for rapid application of measured volumes of toxicants, 261
device to evaluate repellent effect of diphenylmethane, 389
drop net device to determine mosquito prevalence, 321
for irradiation studies, 575
for recording temperatures and humidity of microenvironment, 397
modified mechanical counter, 112
EQ-335, for control of screw-worms, 408
Ereynetes sp., 366
Eriopis convexa, 637
Eristalis arvorum, 147
Erycideus sp., 636
Erythraeus sp., 366
Erythrothrips arizonae, 450
Estigmene acrea, 237, 296, 562
Ethion, for control of
apple grain aphid, 355
codling moth, 357, 379, 422, 542
desert spider mite, 646
European red mite, 355, 422
greenbug, 722
onion maggot, 885
red-banded leaf roller, 379
rosy apple aphid, 355
serpentine leaf miner, 358
soft scales on walnut, 430
two-spotted spider mite, 356, 710, 714, 736
walnut scale, 530
Ethylene chlorobromide, relation between load and concentration of fumigant, 896
Ethylene dibromide
dosages needed to kill rice weevil, 157
relation between load and concentration of fumigant, 896
Ethylene dichloride
dosages needed to kill rice weevil, 157
emulsion, for control of, peach tree borer, 558
Eucelatoria armigera, 562
Euetheola rugiceps, 631
Euschistus tristigmus, 81
Euzesta quadrivittata, 147
Euzophora ostricolorella, 251
Evergestis
pallidata, 281
rimosalis, 254
Exeristes roborator, 652
Exochomus flavipes, 92

Fabric pests, feeding of on natural and synthetic textiles, 248
Fannia
canicularis, 611
spp., 599
Fenson, for control of European red mite, 713
Fenusa pusilla, 245
Flavor evaluations of cacao from trees treated with systemic insecticides, 778
Fleas, on antelope ground squirrel, 32
Fleck, chlorotic, an eriophyid mite injury of plum, 335
Flies, tachinid, reared from lepidopterous larvae in Arizona, 561
Folidol, for control of paddy borer, 411, 571
Forester, or entomologist, 264
Foxella ignota utahensis, 33
Frankliniella
exigua, 394, 450
fusca, 115, 161, 394
gossypiana, 450
insularis, 450
minuta, 450
occidentalis, 450
paucispinosa, 636
sp., 489, 872
tenuicornis, 394
tritici, 115, 161, 394, 451
Fumigants, susceptibility of stages of rice weevil to, 157
Fumigation
effects of methyl bromide on viability of seeds of grain, 847
food commodities for insect control, factors affecting, 891
laboratory tests of organic compounds, 638
methyl bromide against European chafer, 869
methyl bromide, fruit under plastic tarpaulins, 436
warehouse, flue-cured tobacco with HCN against cigarette beetle, 417
Fungi, entomogenous, to control aphids on potato in Maine, 184

Galerucella
notulata, 81
xanthomelaena, 296
Gargaphia torresi, 636
Garmania bulbicola, 366
Gelis tenellus, 650
General Chemical Compound 3707, for control of, house flies, 908
Genite, for control of European red mite, 713
Geocoris
decoratus, 613, 634
punctipes, 757
ventralis, 637
Glycine-sodium hydroxide solution, an attractant for walnut husk fly, 532, 687
Glyodin, in spray combinations against apple insects, 378
Goldfish, toxicity of Sevin to, 540
Goniocotes gallinae, 188
Graphocephala versuta, 81
Grapholitha molesta, 81, 82
Gulfspray naphtha, for control of black carpet beetle, 249
Gummosis in peaches, 27
Guthion, effectiveness against California red scale, 912
Guthion, effect of climatic factors on toxicity of to
boll weevil, 670
desert spider mite, 670
Guthion, effect of on cocooning codling moth larvae, 672
Guthion, for control
boll weevil, 42, 238, 490, 784
bollworm, 42, 238, 490, 646
cabbage leaf-feeding lepidoptera, 443
cabbage looper, 254
codling moth, 379, 422
cotton aphid, 238
cotton fleahopper, 490
cotton leafworm, 259, 784
cross-striped cabbageworm, 254
desert spider mite, 646
European red mite, 422, 714
green peach aphid, 139
house flies, 908
imported cabbageworm, 254
Lithocolletis sp., 423
onion maggot, 885
paddy borer, 411
pink bollworm, 238, 673
red-banded leaf roller, 379
serpentine leaf miner, 358
soft scales on walnut, 430
sugarcane beetle, 633
thrips, 395, 489
two-spotted spider mite, 238, 714, 736
walnut scale, 530
western tussock moth, 764
Guthion, half-life of on cotton leaves, 831

Guthion, toxicity of to
honey bees, 103
pink bollworm adults, 568
Gypona octolineata, 81

Haemagogus equinus, 744
Haematopinus suis, 255
Haplothrips
fasciculatus, 451
purpuratus, 451
Harpalus cautus, 543
HCN, for control of
cigarette beetle, 417
tobacco moth, 417
Health hazards involved in the use of organic phosphorus insecticides, 831
Heat treatment to eliminate European chafer, 912
Heliothis
gelotopoen, 635
spp., 560, 562, 633
virescens, 635, 560
zea, 24, 41, 236, 258, 296, 503, 560, 637, 645, 735, 783, 822
Heliothis zea, effect of fertilizer applications on populations of, 757
Heliothrips haemorrhoidalis, 386
Hemerocampa vetusta, 762
Hemiberlesia
lataniae, 114
rapax, 114
Heptachlor, as a
fumigant against house flies, 638
seed treatment against wireworms, 294
Heptachlor, conversion to epoxide,
in soil, 1
on plants, 3
Heptachlor dips to reduce weevil damage to pine seedlings, 916
Heptachlor, for control of
alfalfa snout beetle, 682
boll weevil, 42, 114, 784
bollworm, 42
cabbage leaf-feeding lepidoptera, 443
cabbage looper, 113
carrot rust fly, 167
clover leaf weevil, 195
clover root curculio, 555
Colorado potato beetle, 829
corn earworm, 823
European chafer, 766
Nysius raphanus, 251
onion maggot, 884
plum curculio, 131
seed-corn maggot, 919
serpentine leaf miner, 358
sugarcane beetle, 631
thrips on cabbage, 395
weevils on crimson clover, 459
western subterranean termite, 209
wireworms, 464, 691
Heptachlor
metabolism of by house flies, 346
percentage recovered from treated soils, 202
persistence of in soils, 11
residues on alfalfa, 222, 392
Heptachlor, toxicity of to
army cutworm, 726
brown-banded cockroach, 610
pink bollworm adults, 578
Scolytus multistriatus, 257
cabbage and muskmelon plants, 534
Hercothrips
braziliensis, 636
marginipennis, 451
Hercules AC-426, for control of chicken body louse, 230
Hercules AC-528, effectiveness against California red scale, 912
Hercules AC-528, for control of
codling moth, 422
corn earworm, 823
cotton leafworm, 259
desert spider mite, 646
European red mite, 714
lone star tick, 425
Mexican bean beetle, 554
mites attacking eye gnat cultures, 462
onion maggot, 885
screw-worm, 425
stable fly, 425
two-spotted spider mite, 714, 736
western tussock moth, 764
Hercules AC-528
resistance of citrus red mite to, 233
toxicity of to honey bees, 103
Heterobostrychus aequalis, 556
Heterothrips prospopidis, 451
Hexamethylphosphoramide, distribution of radioactive in the cockroach, 49
Hibernation, survival and emergence of boll weevils, 745
Hippelates, 278, 557
bishoppi, 557
collusor, 461
pusio, 557
Hippiscus rugosus, 345
Hippodamia
convergens, 375, 612, 722, 759
parenthesis, 614
sinuata disjuncta, 613
quinquesignata, 375
Holcomb C-326, toxicity to honey bees, 103
Honey bee (see *Apis mellifera*) influence of on production
of crimson clover seed, 523
substitute sugar source for, 287
wintering in Minnesota, 332
Hoplopsyllus anomalus, 33
Horcias nobilellus, 636
Horistonotus, 634
Hyalopterus
atriplicis, 179
pruni, 227
Hydrocyanic acid, dosages needed to kill rice weevil, 157
Hydrogen cyanide, relation between load and concentration of fumigant, 896
Hydrogen phosphide, as a fumigant against grain and flour insects, 900
Hydrolysate, yeast, influence of on melon fly, 140
Hylastinus obscurus, 491
Hylemya
antiqua, 883
brassicae, 534
cilicrura, 690, 704, 919
Hylobius pales, 915
Hylurgopinus rufipes, 789
Hypera
brunneipennis, 296
meles, 459
nigrirostris, 459
postica, 392, 506
punctata, 195
Hyperaspis, 614
festiva, 634, 637
jucunda, 92
lateralis, 90
punctata, 366
sp., near *globula*, 92
sp., 770
Hypoderma
bovis, 359, 445, 545, 582, 793, 808, 876
lineatum, 359, 445, 545, 582, 793, 808, 876
spp., 119

Ichneumonidae, characteristics of in relation to biological control, 650
IN-930 as synergist for pyrethrins, 742
Insects
associated with passion fruit flowers in Hawaii, 147
attracted to light traps, 550
control of, affecting birdsfoot trefoil seed production, 492
control programs, function of fish and Wildlife Service in, 740
effectiveness of Dow ET-57 as a fumigant against, 409
Insect
evaluation of plastic coated glass fiber screens, 864
predators, effect of insecticides on alfalfa, 615
role of weather and entomophagus species in natural control of, 474
survival in jet aircraft, 239
thermal death points of, 344
transient, in Delaware's apple and peach plantings, 81
Insecticides
against paddy borer in Taiwan, 571
broadcaster for applying granules, 278
changes in common names of, 373
decline of in soil, 198
dialkyl aryl phosphorothioate, hydrolysis of the alkylphosphate bond by rats, cockroaches, and alkali, 800
effect of certain organic on mortality and oviposition of pink bollworm, 567
effect of climatic factors on toxicities of, 669
effect of on codling moth larvae seeking cocooning quarters, 672
effect of on hatchability and taste, 190
effect of seed treatments of cotton with systemic, 697
effect of soil drenches on young cabbage and muskmelon plants, 534
effect of temperature on against boll weevil, 93
factors influencing recovery and transformation of in soil, 214
granulated, for control of carrot rust fly, 167
for control of insects on snap beans, 258
granulated for control of weevil on crimson clover, 459
granulated for wireworm control, 463
granulated, systemics against spotted alfalfa aphid, 122
granules of heptachlor against boll weevil, 114
fumigation tests of organic compounds, 638
induce increase of soft scales on walnut, 427
insect predators collected on alfalfa before and after treatment with, 615
movement of in soils, 380
multiple regression to compare results of control of meadow spittlebug, 743
new, for control of codling moth, 422
organic, effect of on parasites of oriental fruit moth, 82
organophosphate, developed on rational principles, 714
organophosphorus, metabolism of by microorganisms, 59

organic phosphorus, public health hazards involved in the use of, 831
organophosphorus, resistance of house flies to, 616
residual effect of for control of European chafer, 765
residual sprays on dairy farms against house flies, 908
residual toxicity of to smaller European bark beetle, 256
residues of Diazinon on cherries, 595
residues of on alfalfa, 392
residues of on vegetables, 818
residues of Thimet in small grains grown in treated soil, 623
seed treatments against wireworms, 690
soil, against plum curculio, 131
stable fly resistance to DDT in Norway, 599
susceptibility of brown-banded cockroach to, 608
susceptibilities of four species of potato-infesting aphids to, 414
systemic, action on bloodsucking ectoparasites of 28 compounds given orally to rabbits, 499
systemic, against green peach aphid, 138
systemic, effect of seed treatments on cotton, 495
systemic, field tests with for control of cattle grubs, 793
systemic, for control of cattle grubs, 582
systemic, for control of *Scolytus multistriatus*, 789
systemic, for protection against spotted alfalfa aphid, 457
systemic, for *Theobroma cacao*, 773
systemic, translocation and persistence of in cacao foliage, 773
systemic, trunk implantation in conifers, 400
Thimet granules against leaf miners, 245
timing applications to control European corn borer, 133
Ionization and insect toxicity, 169
Iridomyrmex humilis, 147, 480
Irradiated wheat, effects of on stored grain insects, 674
Irradiation, visible spectrum, effect of on growth in insects, 573
Iseropus orgyiae, 764
Isobornyl thiocyanoacetate, toxicity of in soil to *Ceratitis capitata*, 235
Isodrin, for control of
alfalfa snout beetle, 683
European chafer, 766
onion maggot, 884
serpentine leaf miner, 358
Isolan, translocation and persistence of in cacao foliage, 773
Isomerism, *cis-trans*, effect of on potency of an insect attractant, 921
Iso-Systox, resistance of citrus red mite to, 233
Isovaleryindandione, effect on confused flour beetle, 302
Itonida balsamicola, 400
Itonididae, 366
Itoplectis conquisitor, 650

Jadera sanguinolenta, 635

K.A.A.D. mixture, 740
Karathane, for control of *Aceria tulipae*, 70
Karathane, toxicity of to honey bees, 103
Kelthane, for control of
chicken mite, 911
desert spider mite, 646
European red mite, 255, 714
two-spotted spider mite, 714, 736
Kelthane, resistance to by
Tetranychus atlanticus, 286
Tetranychus pacificus, 286
Kepone, for control of onion maggot, 887
Kerosene, effectiveness against California red scale, 912
Khapra beetle, food habits of larva, 465
Kurtomathrips morrilli, 451

Labauchena daguerrei, 112
Labidura riparia, 366
Lamprophorus tenebrosus, 390
Laphygma
exigua, 296, 562
frugiperda, 562, 634
Lasioderma serricorne, 417
Lasioseius penicilliger, 770
Lasius alienus americanus, 555
Lead arsenate, for control of
bagworm, 368
cherry fruit flies, 594
codling moth, 357, 379, 542
red-banded leaf roller, 379
serpentine leaf miner, 358
Lecanium
cerasorum, 427, 527
corni, 427
pruinosum, 427
Lecopis bella, 90
Ledermuelleria n. sp., 770
Leioseius minusculus, 767
Lepidosaphes beckii, 476
Leptinotarsa decemlineata, 345, 828
Leptocorixa acuta, 405
Leptomastidea abnormis, 92
Leptomastix dactylopii, 92
Leptoterna dolabratus, injury to peaches, 28
Leptothrips
sp. near *mali*, 451
sp., 636
Lerodea eufala, 562
Lethane 384, for control of
black carpet beetle, 249
horn flies, 73, 154
stable flies, 73, 154
Lethane 384 in treatments against biting flies on cattle, 620
Lettuce-yellows prevention, with reference to aster-yellows virus infection and transmission by *Macrosteles fascifrons*, 512
Leucocytozoon disease, effect of black fly control on the incidence of, 845
Leucophenga varia, 733
Light, black, response of hornworm moths to, 78
Light traps, insects attracted to, 550
Limonius agonus, 690
Lindane, as a
fumigant against house flies, 638
seed treatment against wireworms, 294
Lindane, for control of
Aceria tulipae, 70
alfalfa snout beetle, 683
bed bug, 242
black carpet beetle, 249
body lice, 742
carrot rust fly, 166
Ceratitis capitata, 235
chicken body louse, 229
clover leaf weevil, 195
cyclamen mite, 432
horn flies, 559
house flies, 908
melon fly, 236
plum curculio, 132
Pagiocerus fiorii, 468
Scolytus multistriatus, 257
serpentine leaf miner, 358
tropical rat mite, 413
western subterranean termite, 207
wireworms, 691
Lindane
effectiveness of for protection of stored wheat, 843
effect of butter fat content of milk on toxicity to adult house flies, 561
movement of in soils, 381
oral chemotherapeutic activity of in mice, 127
relative toxicity of brown-banded cockroach to, 610
residues of in milling fractions from treated wheat, 516
residues of on vegetables, 818
resistance of house fly to, 497
to prevent emergence of smaller European elm bark beetle, 470
Lindane, toxicity of to
buckthorn aphid, 414
foxglove aphid, 414
green peach aphid, 414
potato aphid, 414
Theriotaphis maculata and *Hippodamia*, 377
Lindorus lophanthae, 479
Linognathus vituli, 877
Lipeurus caponis, 188
Liriomyza pusilla, 357
Liroaspes sp., 366
Listronotus oregonensis, 37
Lithocolletis sp., 423
Lixophaga diatraea, 163
Lobodiplosis sp., 366
Loxostege similalis, 562
Lures, see baits
Lyctocoris campestris, 764
Lygus
lineolaris, 28, 81, 493, 729
pallidulus, 636
Lysiphelbus spp., 375

Macrocentrus
ancylivorus, 85
delicatus, 86
Macronoctua onusta, 549
Macrosiphum
albifrons, 179
creelii, 179
granarium, 178, 553
nigromaculosum, 179
pisi, 169, 178, 613
solanifolii, 144, 184, 414
Macrosteles fascifrons, 512, 903
Mansonia perturbans, 320
Marietta
graminicola, 363
mexicana, 115
picta, 365
Malathion, effect of on soft scales on walnut, 429
Malathion, for control of
Aceria tulipae, 70
bagworm, 368
bed bug, 241
black carpet beetle, 249
body lice, 742
boll weevil, 42, 96, 490, 784
bollworm, 42, 490
cabbage leaf-feeding lepidoptera, 443
California red scale, 257
cherry fruit flies, 594
chicken body louse, 229

chicken mites, 158
citrus cutworm, 656
citrus red mite, 257
citrus whitefly, 552
clover leaf weevil, 196
codling moth, 379
corn earworm, 259, 823
cotton fleahopper, 490
cotton leafworm, 259, 784
cyclamen mite, 432
Epilachna vigintioctopunctata, 749
Fannia canicularis, 611
Fiorinia externa, 406
garden tortrix, 597
hog louse, 255
holly bud moth, 88
honey bees, 108
horn flies, 559
house fly, 361, 908
imported cabbageworm, 254
leaf curl plum aphid, 228
lesser mealworm, 112
loblolly mealybug, 556
Mexican bean beetle, 259, 554
northern fowl mite, 158
Nysius raphanus, 251
onion maggot, 885
Pagiocerus fiorii, 468
potato leafhopper, 259
red-banded leafroller, 379
resistant spotted alfalfa aphids, 313
serpentine leaf miner, 358
six-spotted leafhopper, 903
smaller European elm bark beetle, 470
thrips, 395, 489
two-spotted spider mite, 736
walnut scale, 528
western subterranean termite, 209
western tussock moth, 764

Malathion
detection of by potato flea beetle, 825
effective for protection of stored wheat, 843
ineffective against variegated cutworm, 410
oral chemotherapeutic activity of in mice, 127
suspected cases of poisoning in the Mississippi Delta area, 832

Malathion, residues of on
alfalfa, 392
milling fractions from treated wheat, 516
turnip tops, collards, snap beans, and lettuce, 307
vegetables, 818

Malathion, resistance to by
citrus red mite, 233
house flies, 18, 452, 497, 546, 616

Malathion, toxicity of to
boll weevil, 670
buckthorn aphid, 414
Ceratitis capitata, 235
foxglove aphid, 414
green peach aphid, 414
melon fly, 236
internal parasites of spotted alfalfa aphid, 375
pink bollworm adults, 578
potato aphid, 414
Scolytus multistriatus, 257

Malathion, toxicity and anticholinesterase activity compared with acethion, 716
Medetera aldrichii, 604
Megachile sp., 466
Melanoplus
femur-rubrum, 345
spp., 866
Melanotus
communis, 291, 463
sp., 690
Melcha ornatipennis, 390
Melophagus ovinus, 585
Menacanthus stramineus, 229
Menopon gallinae, 188, 558
Mesostigmata, 259, 768
Metacide, effect of on parasites of oriental fruit moth, 83
Metacide for control of
cyclamen mite, 431
serpentine leaf miner, 358
Metagonistylum minense, 163
Metaphycus
helvolus, 247
lounsburyi, 247
luteolus, 274
stanleyi, 247

Methods
application of insecticides against carrot rust fly, 165
artificial dissemination of fungi against spotted alfalfa aphid, 341
automation system for insectaries, 109
colorimetric whole-blood, to measure cholinesterase activity, 326
determining persistence of heptachlor in soils, 12
evaluating insecticide-treated soils against termites, 207
evaluating sweetclover weevils' preference for sweetclover plants, 190
field studies of the balsam woolly aphid, 540
for proving conversion of, heptachlor to its epoxide on plants, 3; aldrin to dieldrin on plants, 8
for studying biology of carrot weevil, 37
for studying decline of soil insecticides, 199
killing and fixing lepidopterous larvae, 740
polyethylene cover to suppress house fly breeding, 411
probit analysis on a digital computer, 738
radioactive tracer, to study responses of wireworms to chemical seed treatments, 291
rapid determination of sex in walnut husk fly, 551
rapid rearing of clothes moth eggs, 919
rearing carpenterworms, 406
rearing clover root curculio larvae for research uses, 917
rearing large colonies of *Aceria tulipae* in pure culture, 303
sampling to determine prevalence of *Mansonia perturbans*, 320
screening and evaluating compounds for chemotherapeutic activity, 126
self-applicating devices in horn fly control, 795
studying absorption and metabolism of heptachlor by house flies, 346
studying microenvironment of the spotted alfalfa aphid, 397
synthesis of carboxy-labeled C^{14}-*N*,*N*-diethyl-*m*-toluamide, 707
timing of applications against European corn borer, 133
treadle sprayer and cable-type back rubber to control biting flies, 149
ultraviolet determination of diethyltoluamide, 694
use of brine shrimp for bioassay, 780
use of vermiculite in medium for stable fly larvae, 250
wintering honey bees in Minnesota, 332

Methoxychlor, for control of
bed bug, 242
biting flies on cattle, 618
black carpet beetle, 249
cabbage leaf-feeding lepidoptera, 443
cherry fruit flies, 594
chicken body louse, 230
clover leaf weevil, 196
codling moth, 542
holly bud moth, 88
horn fly, 149, 276, 559, 746, 795
house flies, 908
imported cabbageworm, 254
Pagiocerus fiorii, 468
serpentine leaf miner, 358
stable fly, 149, 276
thrips on cabbage, 395
tropical rat mite, 413

Methoxychlor
oral chemotherapeutic activity of in mice, 127
residues of in milling fractions from treated wheat, 516
residues of on vegetables, 818
resistance of house fly to, 497

Methyl bromide
dosages needed to kill rice weevil, 157
effect of different food commodities on concentration of fumigant, 895
effects of fumigation on viability of seeds of grain, 847
fumigation against European chafer, 869
fumigation of fruit under plastic tarpaulins, 436

Methyl chlorobenzilate, toxicity of to honey bees, 103
Methyl demeton, for control of
Aceria tulipae, 70
serpentine leaf miner, 359

Methyl demeton
resistance of citrus red mite to, 233
translocation and persistence of in cacao foliage, 773

Methyl parathion, for control of
boll weevil, 42, 490, 670, 784
bollworm, 42, 490
cabbage looper, 113
cotton fleahopper, 490
cotton leafworm, 784
desert spider mite, 671
garden tortrix, 597
greenbug, 722
Nysius raphanus, 251
onion maggot, 885
paddy borer, 411
thrips, 489
western tussock moth, 764

Methyl parathion
half-life of on cotton leaves, 831
hydrolysis of the alkylphosphate bond by rats, cockroaches, and alkali, 800

Metopius, 652
Metriona bicolor, 81
MGK-264, for control of
horn flies, 73, 276
stable flies, 73, 270, 276
Microcephalothrips abdominalis, 451
Microgaster russata, 390
Milichiella lacteipennis, 147
Mites
parasitic on Alabama mammals, 260
soil-inhabiting from Connecticut apple orchards, 767
Mocis sp., 636
Moisture, effect of on development of weevils and fungi in wheat, 701

Molecular configuration, effect of on toxicity to house flies, 181
Monarthropalpus buxi, 245
Monocephora bicincta, 81
Monocrepidius vespertinus, 81
Monohydrated copper sulfate, toxicity of to honey bees, 103
Monsanto CP-7769, for control of
boll weevil, 490, 785
bollworm, 490
cotton fleahopper, 490
desert spider mite, 646
thrips, 490
Monsanto 8574, for control of two-spotted spider mite, 736
Monsanto 10502, for control of two-spotted spider mite, 736
Monsanto 10851, for control of two-spotted spider mite, 736
Monsanto 11327, for control of two-spotted spider mite, 736
Monsanto 11448, for control of two-spotted spider mite, 736
Moranila californica, 247
Musca domestica, 1, 18, 39, 43, 76, 169, 181, 211, 239, 250, 345, 346, 361, 384, 386, 409, 411, 413, 452, 546, 559, 560, 579, 599, 612, 616, 638, 647, 666, 675, 715, 742, 746; 750, 791, 798, 810, 839, 880, 908, 914
Muscina stabulans, 612
Myzocallus pulchellus, 179
Myzus
monardae, 179
persicae, 138, 145, 178, 414, 812
solani, 145, 414

Nabam in seed treatments of cotton, 697
Nabis
alternatus, 613
spp., 757
Narlene, for control of human bot fly, 724
Naupactus sp., 636
Neisthrea pictipes, 636
Nemorilla floralis, 562
Neoclytus acuminatus, 81
Neodiprion
lecontei, 400
nigroscutum, 125
swainei, 125
Neolygus, damage to peaches, 28
canadensis, 28
caryae, 29
omnivagus, 29
geneseensis, 28
quercalbae, 29
Neotran, persistence of residues, 914
Nephus sp., 92
Netelia, 652
Nezara viridula, 471
Niagara 908, for control of two-spotted spider mite, 736
Niagara 1240 (see Nialate)
Nialate, for control of leaf curl plum aphid, 229
Nicotine, for control of
leaf curl plum aphid, 228
serpentine leaf miner, 358
walnut scale, 528
Nicotine, toxicity of to
Ceratitis capitata, 235
Therioaphis maculata and *Hippodamia*, 377
N,N-diethyl-*m*-toluamide, an insect repellent, 707
N,N-diethyl-*m*-toluamide, for control of horn flies, 276
stable flies, 276
Norvellina seminuda, 81
Nosema disease, 46
NPD, for control of
Aceria tulipae, 70
serpentine leaf miner, 358
Nutrition, comparative NPK, of mites on apple trees, 369
Nysius
raphanus, 250
stimulans, 636

Obituary
Orlando S. Bare, 565
Dwight Fletcher Barnes, 117
Joseph M. Davis, 415
Henry James Franklin, 564
John V. Schaffner, Jr., 266
Harry Scott Smith, 266
Mathis Charles Tanner, 117
Oils, angelica, for trapping Mediterranean fruit flies, 681
Oils, California spray, for control of
California red scale, 193
citrus red mite, 193
Oil, dormant, for control of European red mite, 713
Oil, effect of on transpiration of citrus, 317
Oil emulsion, for control of
citrus whitefly, 552
soft scales on walnut, 427, 429
Oils, for control of
leaf curl plum aphid, 228
rufous scale, 404
Oil, miscible, for control of European red mite, 355
Oil, petroleum, for control of citrus bud mite, 659
Oil-base sprays, for control of black carpet beetle, 249
Olla n. sp., 637
Ololaelaps
hemisphaera, 771
placentula, 771
Oncometopia undata, 81
Oncopeltis fasciatus, 169, 573
Ophionini, 652
Ophyra nigra, 147
Orasema sp., 112
Orchopeas leucopus, 33
Oribatei, 769
Orius
insidiosus, 390, 757
sp., 637
tristicolor, 613
Ornithonyssus
bacoti, 413
sylviarum, 158, 188
Orthopodomyia
alba, 744
signifera, 744
Orthorix, for control of *Aceria tulipae*, 70
Oryzaephilus surinamensis, 674, 900
Otosema odorata, 637
Ovex, for control of
European red mite, 355, 714
mites, attacking eye gnat cultures, 462
two-spotted spider mite, 714
Oxycarenus hyalipennis, 633

Pachylobius picivorus, 915
Pagiocerus fiorii, 468
Panonychus
citri, 193, 232, 353, 914
ulmi, 255, 263, 352, 356, 369, 422, 527, 713
Pantomorus
godmani, 284
sp., 636
Paracalocoris scrupeus, injury to peaches, 28
Paraphaenodiscus sp., 366
Paraphlepsius irroratus, 81
Parascaptomyza disticha, 733
Parasites
attempts to establish against sugarcane borer in La., 163
effect of organic insecticides on, 82
release of in California against citrus mealybugs, 90
species of *Thysanus*, 114
Parasitoidea, 767
Parasitus americanus, 767
Parathion, effect of on
cocooning codling moth larvae, 672
insect predators on alfalfa, 615
parasites of oriental fruit moth, 83
soft scales on walnut, 429
Parathion, for control of
Aceria tulipae, 70
alfalfa pests, 730
alfalfa snout beetle, 683
bagworm, 368
bollworm, 646
cabbage leaf-feeding lepidoptera, 443
carrot rust fly, 166
California red scale, 257, 912
citrus cutworm, 656
citrus red mite, 257
codling moth, 357, 423
corn earworm, 259, 823
cotton leafworm, 758
cyclamen mite, 431
desert spider mite, 646
European chafer, 766
Fannia canicularis, 611
garden tortrix, 597
greenbug, 722
green peach aphid, 139
holly bud moth, 88
house flies, 20, 908
leaf plum aphid, 228
Mexican bean beetle, 259
onion maggot, 885
peach tree borer, 558
potato leafhopper, 259
red-banded leafroller, 456
resistant spotted alfalfa aphids, 313
rufous scale, 404
serpentine leaf miner, 358
six-spotted leafhopper, 903
soft scales on walnut, 427, 429
thrips on cabbage, 395
two-spotted spider mite, 711, 736
walnut aphid, 430
walnut scale, 528
western subterranean termite, 209
western tussock moth, 764
wireworms, 691
Parathion
hydrolysis of the alkylphosphate bond by rats, cockroaches and alkali, 800
movement of in soils, 382
oral chemotherapeutic activity of in mice, 127
not oxidized in *Chlorella* culture, 59
suspected cases of poisoning in the Mississippi Delta area, 832
susceptibility of to brine shrimp, 780
Parathion, resistance to by
citrus red mite, 233
house fly, 497, 546, 616
Tetranychus atlanticus, 286
Tetranychus cinnabarinus, 285
Tetranychus pacificus, 286
Parathion, toxicity of to
Ceratitis capitata, 235
desert spider mite, 670
five species of tetranychid mites, 521
honey bees, 108

internal parasites of spotted alfalfa aphid, 375
melon fly, 236
pink bollworm adults, 578
Pathogens for control of tobacco insects, 560
Pathology, gross, of American cockroach, 663
Pauridea peregrina, 92
Pectinophora gossypiella, 236, 309, 567, 633, 673
Pediculus humanus humanus, 484, 665, 741, 742
Pemphigus
monophagus, 179
populi-caulis, 179
Pentachlorophenyl, for control of western subterranean termite, 207
Pentatomidae, injury to peaches, 27
Perezia pyraustae, 390
Pergamasus
crassipes, var. *longicornis*, 770
sp. 769
Peridroma margaritosa, 410
Periplaneta americana, 49, 169, 211, 239, 409, 574, 663, 715, 866
Perthane, for control of
black carpet beetle, 249
cabbage leaf-feeding lepidoptera, 443
cherry fruit flies, 594
codling moth, 542
garden tortrix, 597
horn flies, 73
imported cabbageworm, 254
onion maggot, 886
serpentine leaf miner, 358
stable flies, 73
thrips on cabbage, 396
Petroleum distillate in treatments against biting flies on cattle, 620
Phaedonia inclusa, 390
Pheidole megacephala, 147
Phenothiazine, used with systemic insecticides, 794
Philaenus lencophthalmus, 81, 219, 492, 743
Phobocampe sp., 764
Phoridae, 112
Phormia regina, 407
Phorocera
fuscimacula, 764
tachinomoides, 562
Phorodon menthae, 179
Phosdrin, for control of
Aceria tulipae, 70
boll weevil, 670
cabbage leaf-feeding lepidoptera, 443
cabbage looper, 254
codling moth, 379
corn earworm, 823
cotton leafworm, 259
cross-striped cabbageworm, 254
desert spider mite, 670
imported cabbageworm, 254
iris borer, 550
Nysius raphanus, 251
onion maggot, 885
red-banded leafroller, 379
resistant spotted alfalfa aphids, 313
serpentine leaf miner, 358
thrips on cabbage, 396
two-spotted spider mite, 736
western tussock moth, 764
Phosdrin
resistance of citrus red mite to, 233
translocation and persistence of in cacao foliage, 773
Phosdrin, toxicity of to
honey bees, 103
Therioaphis maculata and *Hippodamia*, 377
Phosphatase activity of susceptible and resistant flies and cockroaches, 211
Phosphoramides, chromatographic behavior, partition distribution, and anticholinesterase activity of, 52
Phosphorodithioates, carbamoyl alkyl, as chemotherapeutic agents, 126
Phostex, for control of
leaf curl plum aphid, 229
soft scales on walnut, 429
two-spotted spider mite, 736
Phostex, toxicity of to
honey bees, 103
five species of tetranychid mites, 521
Photography, insect, aids in field, 439
Phyllocharis undulata, 390
Phytocoris, injury in peaches, 28
infuscatus, 28
puella, 28
salicis, 28
Phytomyza ilicis, 245
Phytoseiidae, 366, 770
Pieris rapae, 254, 394, 443, 626
Piezodorus quildinii, 636
Piperonyl butoxide, 76
as synergist for pyrethrins, 742
in treatments against biting flies on cattle, 620
synergistic with rotenone against body lice, 484
toxicity and synergism of mixtures with *cis* isomers of allethrin, 675
Pival in bait against confused flour beetle, 301
Plagiodera versicolora, 390
Plagiognathus spp., 493
Plagiomima cognata, 562
Plants, toxicant producing, introduced into Indonesia, 390
Platencyrtus parkeri, 366
Plathypena scabra, 735
Platynota stultana, 296, 598
Plutella maculipennis, 254, 390, 394, 443
Podisus, 637
Polistes exclamans arizonensis, 465
Pollination
birdsfoot trefoil by honey bees, 494
influence on production of crimson clover seed, 523
low-bush blueberries by honey bees, 544
of passion fruit in Hawaii, 146
Polyplax sp., 130
Popillia japonica, 81, 247, 345
Populations
of citrus bud mite, factors affecting, 657
plum curculio in unsprayed peach orchard, 330
Powder-post beetles, Bostrichid, host records of, 556
Praon palitans, 374
Predators, release of in California against citrus mealybugs, 90
Prionoxystus robiniae, 407
Prodenia
eridania, 637, 838
ornithogalli, 562
sp., 390
Prospaltella spp., 406
Protoparce
quinquemaculata, 78, 560
sexta, 78, 560
Protozoa, disease-producing in *Trogoderma*, 916
Psallus seriatus, 41, 489, 872
Pseudaletia unipuncta, 562
Pseudaphycus
angelicus, 92
perdignus, 92
Pseudococcus
adonidum, 90, 433
brevipes, 905
citri, 90, 480, 773
gahani, 90
maritimus, 90
njalensis, 773
Pseudomonas fluorescens, 59
Pseudopeziza, 220
PS-256, tests against northern fowl mite, 188
Psila rosae, 165
Psilothrips
paradalotus, 451
priesneri, 451
Pterostichus vulgaris, 543
Ptilodactyla serricollis, 81
Pulex irritans, 33
Pyemotidae, 461
Pyrausta nubilalis, 133, 735, 744
Pyrazoxon, resistance of citrus red mite to, 233
Pyrenone, for control of
horn flies, 73
Pagiocerus fiorii, 468
stable flies, 73
Pyrenone, toxicity of in soil to *Ceratitis capitata*, 235
Pyrethrins, for control of
black carpet beetle, 249
body lice, 742
horn flies, 73, 154, 276
Nysius raphanus, 251
stable flies, 73, 154, 270, 276
Pyrethrins
in treatments against biting flies on cattle, 620
synergized, for control of bed bug, 241
toxicity of in house fly sprays, 810
with acetal synergists in house fly sprays, 39
Pyrethrum, toxicity of to *Therioaphis maculata*, and *Hippodamia*, 377
Pyrolan, toxicity of in soil to *Ceratitis capitata*, 235

Radioactive materials
absorption and toxicity in mosquito larvae, 640
carboxy-labeled C^{14}-*N*,*N*-diethyl-*m*-toluamide, 707
cobalt-60 to tag wireworms, 291
dimefox in the cockroach, 50
fate of P^{32}-labeled Bayer 21/199 in the white rat, 204
P^{32} parathion in soils, 380
schradan in the cockroach, 50
(S^{35}) Bayer 19639 in pineapple plants, 905
Raphignathus sp., 366
Rearing, laboratory of the boll weevil, larval diet, 288
Red clover, yield losses caused by clover root borer, 491
Reduviidae, 549
Regression, multiple, to compare several meadow spittlebug insecticides, 743
Repellents
against horn flies and stable flies on beef cattle, 276
insect, synthesis of, 707
insect, ultraviolet determination of diethyltoluamide, 694
insecticide seed treatments against wireworms, 291

Residues
in swine sprayed with malathion, 256
persistence of aldrin on alfalfa, 222
persistence of heptachlor on alfalfa, 222
Residues of
Chipman R-6200 in cacao, 776
commonly used insecticides on alfalfa, 392
DDT and malathion on turnip tops, collards, snap beans, and lettuce, 306
demeton in cacao, 777
demeton on cabbage and celery, 175
dieldrin on alfalfa treated with granules, 685
dieldrin in aldrin-treated carrots, 340
insecticide, in milling fractions in wheat treated with insecticides, 516
insecticide, in soils, 186
insecticides on vegetables, 818
Isolan in cacao, 777
lindane on and in treated wheat, 844
malathion on and in treated wheat, 844
methoxychlor in milk of treated cattle, 618
Neotran on and in lemons and oranges, 914
Thimet in cacao, 777
Thimet in small grains grown in treated soil, 623
toxic non-aldrin in carrots grown on aldrin-treated soil, 337
Residues, bromine
in avocados after fumigation with ethylene chlorobromide and ethylene dibromide, 899
in commodities after fumigation with methyl bromide, 896
Resistance
alfalfa to meadow spittlebug, 219
alfalfa to spotted alfalfa aphid, 457, 760
bed bugs to DDT, 241
boll weevil to toxaphene, 721
citrus red mite to organic phosphates, 232
codling moth to DDT, 693
Colorado potato beetle to DDT, 828
Culex tarsalis to DDT, 56
DDT, attempt to reduce in body lice, 665
difference in damage to varieties of sweet corn by Japanese beetle, 247
effects of chilling pupae on emergence of DDT-resistant house flies, 666
effect of seeding density of spring wheat on wheat stem sawfly, 804
effect of temperature on insecticides against boll weevil, 93
European red mite to phosphorus-based acaricides, 255
genotypic reactions to boll weevil attack in Upland cotton, 508
potato flea beetle to DDT, 823
reproduction in insecticide-treated house flies, 579
spotted alfalfa aphid to organophosphorus insecticides, 312
stable flies to DDT in Norway, 599
sweetclover to sweetclover weevil, 190
Tetranychus to organophosphorous insecticides, 285
to boll weevil in mixed populations of resistant and susceptible cotton plants, 502
winter wheat to the greenbug, 702
Resistance of house fly to
Bayer 21/199, 497
chlordane, 497
dicapthon, 497
DDT, 497, 647
Diazinon, 497, 546, 616
dieldrin, 497
Dipterex, 497, 616
Dow ET-57, 497
lindane, 497
malathion, 18, 452, 497, 546, 616
methoxychlor, 497
organophosphorus compounds, 20, 616
parathion, 497, 546, 616
Reticulitermes hesperus, 207
Rhadinopsylla heiseri, 33
Rhagoletis
completa, 532, 551, 686
cingulata, 592
fausta, 592
Rhopalosiphum
fitchii, 354, 553
maidis, 185, 553, 637
nymphaeae, 227
padi, 179
poae, 179
Rhopobota naevana, control of, 87
Rhyacionia frustrana, 556
Rhyssa persuasoria, 650
Rhyzopertha dominica, 843, 900
Rotenone
for control of Mexican bean beetle, 554
ineffective against variegated cutworm, 410
toxicity of to *Therioaphis maculata* and *Hippodamia*, 377
R-326, for control of stable flies, 270
Ryania, for control of
codling moth, 379, 422
Lithocolletis sp., 423
red-banded leafroller, 379
sugarcane borer, 562

Sabadilla, toxicity of to honey bees, 104
Sabulodes caberata, 296
Safety precautions violated in the use of insecticides, 834
Saissetia oleae, 247
Sanninoidea exitiosa, 557
Sarcophaga, sp., 634, 637
Scambus
boulianae, 650
nucum, 650
Scarabaeid beetles, human ear invasions by, 546
Schistocerca
americana, 345
paranesis, 633
sp., 465
Schizonotus latus, 390
Schoenobius
incertulas, 411, 571
officinale, 390
Schradan
distribution of radioactive, in cockroach, 49
effect of on soft scales on walnut, 429
Schradan, for control of
Aceria tulipae, 70
European red mite, 713
green peach aphid, 139
Nysius raphanus, 251
serpentine leaf miner, 358
walnut aphid, 527
Schradan not oxidized in *Chlorella* cultures, 59
Schradan, resistance of citrus red mite to, 233
Schradan tends to induce increased populations of
calico scale, 527
European red mite, 527
Schradan, toxicity of to *Therioaphis maculata* and *Hippodamis*, 377
Scirtothrips
citri, 451
sp., 451
Scirpophaga
innotata, 390
nivella, 390
Scobicia declivis, 556
Scolothrips sexmaculatus, 451
Scolytus multistriatus, 256, 469, 789
Screens, plastic coated glass fiber insect, 864
Scutellista cyanea, 247
Scymnus
quadrivittatus, 92
sordidus, 90
sp., 637
Seed treatments, cotton with systemic insecticides, 160
Sehirus cinctus, 81
Selenaspidus articulatus, 404
Sesamia inferens, 571
Sevin, effect of on cocooning codling moth larvae, 672
Sevin, for control of
apple aphid, 422
boll weevil, 236, 490, 784
bollworm, 236, 490, 646
cabbage leaf-feeding lepidoptera, 443
cabbage looper, 236, 254
codling moth, 379, 422, 542
corn earworm, 259, 823
cross-striped cabbageworm, 254
cotton aphid, 236
cotton fleahopper, 490
cotton leafworm, 237, 259
imported cabbageworm, 254
Lithocolletis sp., 423
Mexican bean beetle, 259, 554
onion maggot, 885
pink bollworm, 236
potato leafhopper, 259
red-banded leafroller, 379
salt-marsh caterpillar, 237
sugarcane beetle, 633
thrips on cabbage, 396
thrips, 489
western tussock moth, 764
Sevin, tests against northern fowl mite, 189
Sevin (C-7744), toxicity of to
goldfish, 540
honey bees, 103
Shell OS-1808
for control of, *Aceria tulipae*, 70
translocation and persistence of in cacao foliage, 773
Shell OS 1836
for control of onion maggot, 885
translocation and persistence of in cacao foliage, 773
Siglure (ENT 21486) as a lure for Mediterranean fruit fly, 679
Simulium
congareenarum, 845
decorum, 846
slossonae, 845
tuberosum, 845
venustum, 845
verecundum, 845
Siphona irritans, 72, 149, 269, 275, 559, 618, 746, 795
Sitona
cyclindricollis, 190
hispidula, 554, 917
Sitophilus
granarius, 674, 701, 843, 900
oryza, 23, 157, 674, 843, 860, 900, 913

sasakii, 913
Sitotroga cerealella, 23, 674
S-MP powder, tests against northern fowl mite, 188
Sodium arsenite
for control of western subterranean termite, 207
spray injury to dairy cattle by, 539
Sodium pentachlorophenate, for control of western subterranean termite, 207
Soil drenches, insecticidal, effect of on growth of young cabbage and muskmelon plants, 534
Solenopsis saevissima, 111
Solenoptes capillatus, 877
Solvents, organic, effects of on American cockroach, 664
Speed of action, comparative study, of some halogen-containing thiol-alkylating agents on resistant house flies, 880
Spicaria rileyii, 560
Spodoptera mauritia, 571
Sprays, barn, residual, effect of on horn flies, 559
Staphylinidae, 770
Steneotarsonemus pallidus, 431
Stenotarsus hispidus, 81
Stenotus binotatus, injury to peaches, 28
Stomoxys calcitrans, 72, 149, 250, 269, 275, 425, 499, 599, 612, 618, 861
Strobane, for control of
Aceria tulipae, 70
black carpet beetle, 249
European chafer, 766
serpentine leaf miner, 358
Strobane, toxicity of in soil to *Ceratitis capitata*, 235
Sulfoxide
as synergist for pyrethrins, 742
synergistic with rotenone against body lice, 484
Sulfur for control of
mites attacking eye gnat cultures, 462
Nysius raphanus, 251
Sulfur, relative toxicity of to five species of tetranychid mites, 521
Sulphenone, for control of northern fowl mite, 189
Supella supellectilium, 608
Sympherobius spp., 90
Synergism
compounds showing, with rotenone against body lice, 484
of mixtures of piperonyl butoxide with *cis* isomers of allethrin, 675
piperonyl butoxide and IN-930 with pyrethrins against house flies, 742
sulfoxide and IN-930 with pyrethrins against body lice, 742
Synergists
ability of two polyfluoro BHC analogs to synergize lindane, 413
compounds related to DDT evaluated as DDT synergists, 647
Syrbula admirabilis, 345
Syrphus americanus, 81
Sytam, for control of iris borer, 550

Tabanus
giganteus, 618
quinquevittata, 275
sp., 72
sulcifrons, 274, 618
Tabutrex, for control of stable flies, 270
Tachinidae, reared from lepidopterous larvae in Arizona, 561
Taeniothrips orionis, 274
TDE (DDD), for control of
citrus cutworm, 656
codling moth, 379
garden tortrix, 597
red-banded leafroller, 379, 456
serpentine leaf miner, 358
western tussock moth, 764
Technique (see also methods)
aids for field photography of insects, 439
testing insecticide deposits with newly hatched codling moth larvae, 547
Tedion, for control of
citrus red mite, 352
European red mite, 353, 713
lone star tick, 425
mites attacking eye gnat cultures, 462
screw-worm, 425
stable fly, 425
two-spotted spider mite, 352, 736
Teleonemia scrupulosa, 390
Temperature, effect of on
efficiency of insecticides against boll weevil, 93
fumigation, 892
house fly emergence, 666
survival of aphids, 144
survival rate of European red mite eggs, 263
Tenebroides mauritanicus, 866
TEPP, for control of
citrus cutworm, 656
serpentine leaf miner, 358
TEPP, toxicity of to *Therioaphis maculata* and *Hippodamia*, 377
TEPP, used with polyhedrosis virus, for control of cabbage looper, 629
Tetracnemus
peregrinus, 92
pretiosus, 92
Tetranychidae, 41, 633, 767
Tetranychus
atlanticus, 285
bimaculatus, 767
cinnabarinus, 225, 518
desertorum, 518, 634, 645, 872
gloveri, 518
lobosus, 518
ludeni, 519
mcdanieli, 588
pacificus, 225
tetarius, 238, 352, 356, 369, 519, 710, 713
tumidus, 518, 669
Tetrastichus
marcovitchi, 401
sp., 366
whitmani, 401
Thanasimus nigriventris, 604
Thanite, in treatments against biting flies on cattle, 620
Thanite, for control of
horn flies, 73
stable flies, 73
Therioaphis (*Pterocallidium*) *maculata*, 122, 179, 287, 312, 341, 374, 392, 397, 457, 553, 573, 601, 612, 760
Thermal constants of seed-corn maggot, 704
Thermobia domestica, 248
Thimet, effect of seed treatments on cotton, 495, 697
Thimet, for control of
bagworm, 368
balsam gall midge, 400
birch leaf miner, 245
boxwood leaf miner, 245
boll weevil, 100
cabbage leaf-feeding lepidoptera, 443
citrus mealybug, 779
citrus whitefly, 552
codling moth, 422
corn earworm, 259
cotton aphid, 120
European red mite, 714
green peach aphid, 139
holly leafminer, 245
leaf curl plum aphid, 226
Mexican bean beetle, 259
onion maggot, 885
potato leafhopper, 259
Scolytus multistriatus, 789
serpentine leaf miner, 358
spotted alfalfa aphid, 122, 457
sugarcane beetle, 633
thrips on cabbage, 395
thrips on cotton, 160
two-spotted spider mite, 714, 736
Thimet, metabolism of by
plants, insects, and mammals, 838
soil microorganisms, 59
Thimet
tests with as cottonseed treatments, 872
to prevent blue dwarf disease in oats, 623
toxicity and stability to hydrolysis by alkali, 839
toxicity of to honey bees, 103
translocation and persistence of in cacao foliage, 773
Thiobacillus thiooxidans, 59
Thiodan, for control of
boll weevil, 42, 490, 784
bollworm, 42, 490, 646
cabbage leaf-feeding lepidoptera, 443
cabbage looper, 113, 254
citrus whitefly, 552
corn earworm, 259, 823
cotton fleahopper, 490
cross-striped cabbageworm, 254
greenbug, 722
imported cabbageworm, 254
lone star tick, 425
Mexican bean beetle, 259, 554
onion maggot, 885
potato leafhopper, 259
screw-worm, 425
serpentine leaf miner, 358
soft scales on walnut, 430
stable fly, 425
sugarcane beetle, 633
thrips, 395, 489
two-spotted spider mite, 736
walnut scale, 530
Thiol alkylating agents, halogen-containing, speed of action on resistant house flies, 880
Thrassis bacchi gladiolis, 32
Thrips
control of on cotton with systemics, 698
effect of on height and weight of seedling cotton, 115
hawaiiensis, 147
of Arizona, 450
tabaci, 394, 451
Thyanta accumulata, 636
Thyridopteryx ephemeraeformis, 367
Thysanus
merceti, 114
thoreauini, 114
Timberlakia europaea, 365
Tineola bisselliella, 248, 919
Tomocera (*Moranila*), 247
Torulopsis utilis, 59
Toxaphene
effect of wind and rain on toxicity of to boll weevil, 670
effect on insect predators on alfalfa, 615

Toxaphene, for control of
Aceria tulipae, 70
alfalfa pests, 730
alfalfa snout beetle, 683
boll weevil, 490, 784
bollworm, 490, 646
cabbage leaf-feeding lepidoptera, 443
cabbage looper, 113, 254
citrus cutworm, 656
clover leaf weevil, 195
Colorado potato beetle, 829
corn earworm, 823
cotton fleahopper, 490
cotton leafworm, 259, 784
cross-striped cabbageworm, 254
cyclamen mite, 432
European chafer, 766
greenbug, 722
horn fly, 795
imported cabbageworm, 254
meadow spittlebug bug, 493
Nysius raphanus, 251
seed-corn maggot, 919
serpentine leaf miner, 358
thrips, 396, 489, 698
two-spotted spider mite, 736
weevils on crimson clover, 459
western subterranean termite, 209
western tussock moth, 764
Toxaphene
residues of on vegetables, 818
residues in treated soil, 187
spray compared with cottonseed treatments with systemics, 873
susceptibility of the boll weevil to, 719
tests against boll weevil, 95
Toxaphene, toxicity of to
buckthorn aphid, 414
foxglove aphid, 414
green peach aphid, 414
pink bollworm adults, 568
potato aphid, 414
Therioaphis maculata and *Hippodamia*, 377
Toxoptera
graminum, 553, 701, 722
muklenbergiae, 179
Toxorhynchites rutilus, 744
Transmission of sugar beet yellow-net virus, latent period phenomena in, 812
Traps
for eradicating Mediterranean fruit fly in Florida, 679
for walnut husk fly, 689
light, insects attracted to, 550
light, mechanical and suction-type for hornworm moths, 78
Triatoma sanguisuga, 549
Tribolium confusum, 239, 299, 345, 385, 409, 674, 860, 900
Trichoplusia ni, 113, 236, 254, 296, 394, 443, 562, 626
Trioxys utilis, 374
Trithion, for control of
California red scale, 912
codling moth, 422, 542
corn earworm, 259
desert spider mite, 646
European red mite, 255, 422, 714
leaf curl plum aphid, 228
lone star tick, 425
Mexican bean beetle, 259
northern fowl mite, 189
onion maggot, 883
paddy borer, 411
potato leafhopper, 259
resistant spotted alfalfa aphids, 313
screw-worm, 425
soft scales on walnut, 430
stable fly, 425
two-spotted spider mite, 710, 714, 736
walnut scale, 530
Trithion, resistance of
citrus red mite to, 232
Tetranychus atlanticus to, 286
Tetranychus pacificus to, 286
Trithion, toxicity of to
honey bees, 103
Therioaphis maculata and *Hippodamia*, 377
Trogoderma
glabrum, 916
granarium, 465, 900, 916
inclusum, 916
parabile, 916
simplex, 916
sternale, 763, 916
teukton, 916
Trogus, 652
Trombidiidae, 771
Trombidiformes, 768
Tropidophryne melvillei, 92
Tyndarichus sp., 366

Udea rubigalis, 296
Urginea maritima, 390
Uropodina, 771

Vanessa cardui, 562
Vasates
cornutus, 337
fockeui, 335
V-C 13, for control of onion maggot, 886
Vectors, yellow-fever, in Dallas, Tex., 743
Veigaia
cerva, 770
nemorensis, 770
Vermiculite
as a carrier for insecticides against wireworms, 463
use of in medium for stable fly larvae, 250
Virus disease
Aceria tulipae as vector of, 68
granulosis, for control of red-banded leaf roller, 454
polyhedrosis, for control of cabbage looper, 626
sugar beet yellow-net, transmission of by green peach aphid, 812
Volucella obesa, 147
Voria ruralis, 562

Warfarin, ineffective against confused flour beetle, 302
Water drinking by larvae of European corn borer, 744
Weather
effect of on toxicities of insecticides, 669
role of in natural control of insect populations, 474
Wheat, development of weevils and fungi in columns of, 701
Winthemia sp., 637

Xanthoencyrtus phragmitis, 365
Xenopsylla cheopis, 239
Xylocopa varipuncta, 146
Xylomyges curialis, 653

Yeast hydrolysate, influence of on melon fly, 140
Yields
corn, reduction of by southwestern corn borer, 747
red clover, effect of distance from honey bee colonies on, 64

Zarhopalus sheldoni, 92
Zeadiatreae grandiosella, 747
Zenillia hyphantriae, 562
Zercoseius sp., 366